

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>				
1. REPORT DATE (DD-MM-YYYY) July 2013		2. REPORT TYPE Viewgraph		3. DATES COVERED (From - To) July 2013- August 2013
4. TITLE AND SUBTITLE Acceleration of PCI and CR algorithms for high-fidelity in-space propulsion modeling (Briefing Charts)			5a. CONTRACT NUMBER In-House	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) R. Martin, H. Le, C. Lederman, J.-L. Cambier			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER Q0AE	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRS 1 Ara Drive. Edwards AFB CA 93524-7013			8. PERFORMING ORGANIZATION REPORT NO.	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB CA 93524-7048			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2013-180	
12. DISTRIBUTION / AVAILABILITY STATEMENT Distribution A: Approved for Public Release; Distribution Unlimited. PA#13398				
13. SUPPLEMENTARY NOTES Viewgraph for the AFOSR Computational Mathematics Program Review, Arlington,VA, 29 July – 1 August 2013.				
14. ABSTRACT We describe enhancements under development for multi-scale methods to be applied to the high-fidelity modeling of spacecraft electric propulsion systems and their environment. Multiple challenges in multi-scale integration, statistical accuracy, and physical modeling are addressed through a variety of innovative numerical methods and mathematical approaches. We emphasize the advances made on a specialized multi-scale time-stepping integrator for finite-Larmor radius particle trajectories, accelerated collisional-radiative non-equilibrium ionization kinetics through Boltzmann equilibrated level groups, and a novel approach to dynamic phase-space reconstruction which can be used to resolve the problems of multi-scale statistics in particle-based kinetic simulations.				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 76
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified		
				19a. NAME OF RESPONSIBLE PERSON Jean-Luc Cambier
				19b. TELEPHONE NO (include area code) 661-525-5649

ACCELERATION OF PIC AND CR ALGORITHMS FOR HIGH FIDELITY IN-SPACE PROPULSION MODELING

Robert Martin¹, Hai Le¹, Carl Lederman¹
PI: Jean-Luc Cambier²

ERC INC.¹, IN-SPACE PROPULSION BRANCH²,
AIR FORCE RESEARCH LABORATORY
EDWARDS AIR FORCE BASE, CA USA

AFOSR Computational Math Annual Review,
Arlington VA,
July 29, 2013



U.S. AIR FORCE



Distribution A: Approved for public release; unlimited distribution
PA (Public Affairs) Clearance Number TBD




- 1 INTRODUCTION
- 2 EFFICIENT ELECTRON PUSH
- 3 ACCELERATED COLLISIONAL-RADIATIVE MODELS
- 4 PARTICLE REMAPPING
- 5 FUTURE WORK



Spacecraft Propulsion Relevant Plasma:

- From hall thrusters to plumes and fluxes on components
- Complex reaction physics i.e. Discharge and Breakdown in FRC
- Relevant Densities often Span 6+ Orders of Magnitude
- Spatial scales of interest span μm - $100m$ range

Electric Propulsion Plumes



Chamber Environment



SPACECRAFT PLASMA MODELING CHALLENGES


Spacecraft Propulsion Relevant Plasma:

- From hall thrusters to plumes and fluxes on components
- Complex reaction physics i.e. Discharge and Breakdown in FRC
- Relevant Densities often Span 6+ Orders of Magnitude
- Spatial scales of interest span μm - $100m$ range

Solution?

Multi-Scale and Multi-Physics
Adaptive Algorithms

Electric Propulsion Plumes



Chamber Environment



Boris Push:


$$v^- = v(t - \Delta t/2) + \frac{q\Delta t}{2m} E(t)$$

$$v' = v^- + \frac{q\Delta t}{2m} v^- \times B(t)$$

$$v^+ = v^- + \frac{\frac{q\Delta t}{2m}}{1 + \left(\frac{q\Delta t}{2m} |B|\right)} v' \times B(t)$$

$$v(t + \Delta t/2) = v^+ + \frac{q\Delta t}{2m} E(t)$$

$$r(t + \Delta t) = r(t) + \Delta t v(t + \Delta t/2)$$





Boris Push:

$$v^- = v(t - \Delta t/2) + \frac{q\Delta t}{2m} E(t)$$


$$v' = v^- + \frac{q\Delta t}{2m} v^- \times B(t)$$

$$v^+ = v^- + \frac{\frac{q\Delta t}{2m}}{1 + \left(\frac{q\Delta t}{2m} |B|\right)} v' \times B(t)$$

$$v(t + \Delta t/2) = v^+ + \frac{q\Delta t}{2m} E(t)$$

$$r(t + \Delta t) = r(t) + \Delta t v(t + \Delta t/2)$$

- Leap-Frog in r & v





Boris Push:

$$v^- = v(t - \Delta t/2) + \frac{q\Delta t}{2m} E(t)$$


$$v' = v^- + \frac{q\Delta t}{2m} v^- \times B(t)$$

$$v^+ = v^- + \frac{\frac{q\Delta t}{2m}}{1 + \left(\frac{q\Delta t}{2m} |B|\right)} v' \times B(t)$$

$$v(t + \Delta t/2) = v^+ + \frac{q\Delta t}{2m} E(t)$$

$$r(t + \Delta t) = r(t) + \Delta t v(t + \Delta t/2)$$

- Leap-Frog in r & v
- Energy Conservation only via Interpolation in Time





Boris Push:

$$v^- = v(t - \Delta t/2) + \frac{q\Delta t}{2m} E(t)$$


$$v' = v^- + \frac{q\Delta t}{2m} v^- \times B(t)$$

$$v^+ = v^- + \frac{\frac{q\Delta t}{2m}}{1 + \left(\frac{q\Delta t}{2m} |B|\right)} v' \times B(t)$$

$$v(t + \Delta t/2) = v^+ + \frac{q\Delta t}{2m} E(t)$$

$$r(t + \Delta t) = r(t) + \Delta t v(t + \Delta t/2)$$

- Leap-Frog in r & v
- Energy Conservation only via Interpolation in Time
- Numerically Stable but Drifts and Fails when $\Delta t > \omega_c/2$





Boris Push:

$$v^- = v(t - \Delta t/2) + \frac{q\Delta t}{2m} E(t)$$


$$v' = v^- + \frac{q\Delta t}{2m} v^- \times B(t)$$

$$v^+ = v^- + \frac{\frac{q\Delta t}{2m}}{1 + \left(\frac{q\Delta t}{2m} |B|\right)} v' \times B(t)$$

$$v(t + \Delta t/2) = v^+ + \frac{q\Delta t}{2m} E(t)$$

$$r(t + \Delta t) = r(t) + \Delta t v(t + \Delta t/2)$$


- Leap-Frog in r & v
- Energy Conservation only via Interpolation in Time
- Numerically Stable but Drifts and Fails when $\Delta t > \omega_c/2$





Gyrokinetic Push:


- Effective for High-B Plasma (i.e. Magnetic-Fusion)
- Assumes $\lambda_c \ll dx$
- Loses Phase Information
- Assumes Phase-Scatter Diffusion Negligible





B-Frame Push:


- Coordinates Rotated and Aligned to *B*-Field





B-Frame Push:


- Coordinates Rotated and Aligned to B -Field
- Motion Decomposed into Rotation and Drift
- Exact Solution in Constant Field





B-Frame Push:

- Coordinates Rotated and Aligned to B -Field
- Motion Decomposed into Rotation and Drift
- Exact Solution in Constant Field
- Phase Preserved Across Many- ω_c






B-Frame Push:

- Coordinates Rotated and Aligned to B -Field
- Motion Decomposed into Rotation and Drift
- Exact Solution in Constant Field
- Phase Preserved Across Many- ω_c


What about Variable Fields?





B-Frame Push in Variable Fields:


- Consistent Solution Recovered with Multiple Steps





B-Frame Push in Variable Fields:

- Consistent Solution Recovered with Multiple Steps
- Convergence Order is Low




Magnetic Mirror Test



B-Frame Push in Variable Fields:

- Consistent Solution Recovered with Multiple Steps
- Convergence Order is Low
- Solution Stalls when $dt > \omega_c$



Magnetic Mirror Test



MULTI-SCALE PARTICLE ADVANCEMENT

B-Frame Push in Variable Fields:

- Consistent Solution Recovered with Multiple Steps
- Convergence Order is Low
- Solution Stalls when $dt > \omega_c$
- Consider the Push in \hat{B} -Aligned Coordinates...
- No Mechanism for $v_{\perp} \rightarrow v_{\parallel}$ in Step?
(No Bounce in Magnetic Mirror?)
- Variable \vec{B} and \vec{E} via Taylor Series?

$$\Delta \vec{v} = \mathbf{D}_0 \cdot \vec{v} + \frac{q\Delta t}{m} \mathbf{D}_1 \cdot \vec{E}$$

$$\Delta \vec{x} = \mathbf{D}_1 \cdot \vec{v} \Delta t + \frac{q\Delta t^2}{m} \mathbf{D}_2 \cdot \vec{E}$$

$$\mathbf{D}_k = \hat{\mathbf{R}}^{-1} \cdot \Delta_k \cdot \hat{\mathbf{R}}$$

$$\Delta \mathbf{v}^R = \underbrace{\begin{pmatrix} -C_0 & S_0 & 0 \\ -S_0 & C_0 & 0 \\ 0 & 0 & 0 \end{pmatrix}}_{\Delta_0} \cdot \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} + \underbrace{\begin{pmatrix} S_1 & C_1 & 0 \\ -C_1 & S_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\Delta_1} \cdot \begin{bmatrix} a_x \\ a_y \\ a_z \end{bmatrix}$$

$$\Delta \mathbf{x}^R = \underbrace{\begin{pmatrix} S_1 & C_1 & 0 \\ -C_1 & S_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\Delta_1} \cdot \begin{bmatrix} v_x \Delta t \\ v_y \Delta t \\ v_z \Delta t \end{bmatrix} + \underbrace{\begin{pmatrix} C_2 & -S_2 & 0 \\ S_2 & C_2 & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}}_{\Delta_2} \cdot \begin{bmatrix} a_x \Delta t^2 \\ a_y \Delta t^2 \\ a_z \Delta t^2 \end{bmatrix}$$



MULTI-SCALE PARTICLE ADVANCEMENT

B-Frame Push in Variable Fields:

- Consistent Solution Recovered with Multiple Steps
- Convergence Order is Low
- Solution Stalls when $dt > \omega_c$
- Consider the Push in \hat{B} -Aligned Coordinates...
- No Mechanism for $v_{\perp} \rightarrow v_{\parallel}$ in Step?
(No Bounce in Magnetic Mirror?)
- Variable \vec{B} and \vec{E} via Taylor Series?

But even \hat{R} depends on \vec{B}
(And is Partially Arbitrary!)

$$\Delta \vec{v} = \mathbf{D}_0 \cdot \vec{v} + \frac{q\Delta t}{m} \mathbf{D}_1 \cdot \vec{E}$$

$$\Delta \vec{x} = \mathbf{D}_1 \cdot \vec{v} \Delta t + \frac{q\Delta t^2}{m} \mathbf{D}_2 \cdot \vec{E}$$

$$\mathbf{D}_k = \hat{\mathbf{R}}^{-1} \cdot \Delta_k \cdot \hat{\mathbf{R}}$$

$$\Delta \mathbf{v}^R = \underbrace{\begin{pmatrix} -C_0 & S_0 & 0 \\ -S_0 & C_0 & 0 \\ 0 & 0 & 0 \end{pmatrix}}_{\Delta_0} \cdot \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} + \underbrace{\begin{pmatrix} S_1 & C_1 & 0 \\ -C_1 & S_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\Delta_1} \cdot \begin{bmatrix} a_x \\ a_y \\ a_z \end{bmatrix}$$

$$\Delta \mathbf{x}^R = \underbrace{\begin{pmatrix} S_1 & C_1 & 0 \\ -C_1 & S_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\Delta_1} \cdot \begin{bmatrix} v_x \Delta t \\ v_y \Delta t \\ v_z \Delta t \end{bmatrix} + \underbrace{\begin{pmatrix} C_2 & -S_2 & 0 \\ S_2 & C_2 & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}}_{\Delta_2} \cdot \begin{bmatrix} a_x \Delta t^2 \\ a_y \Delta t^2 \\ a_z \Delta t^2 \end{bmatrix}$$



B-Frame Push in Variable Fields (cont.):

- Push as Explicit Matrix Operator:
(For now Ignoring \vec{E})

$$\mathbb{X}_+ = \mathbb{M} \cdot \mathbb{X}_0$$

$$\mathbb{X}_+ = \begin{bmatrix} \vec{x}_+ \\ \vec{v}_+ \end{bmatrix} = \begin{bmatrix} \vec{x}_0 + \Delta \vec{x} \\ \vec{v}_0 + \Delta \vec{v} \end{bmatrix} = \underbrace{\begin{bmatrix} \mathbb{I} & \mathbf{D}_1 \\ 0 & \mathbb{I} + \Delta t \mathbf{D}_0 \end{bmatrix}}_{\mathbb{M}} \cdot \begin{bmatrix} \vec{x}_0 \\ \vec{v}_0 \end{bmatrix}$$



B-Frame Push in Variable Fields (cont.):

- Push as Explicit Matrix Operator:
(For now Ignoring \vec{E})
- What's Known about \mathbb{M} :
 - Reversible through Time Inversion...
 - \mathbb{M}^{-1} Must Exist
 - \mathbb{M} is a 1:1 Map of $\mathbb{X}_0 \rightarrow \mathbb{X}_+$
 - \mathbb{M} is Unique (Even if \hat{R} was Not...)
 - If \mathbb{M} is Unique, Elements are Unique
 - \mathbb{M} is Constant if \vec{B} is Constant

$$\mathbb{X}_+ = \mathbb{M} \cdot \mathbb{X}_0$$

$$\mathbb{X}_+ = \begin{bmatrix} \vec{x}_+ \\ \vec{v}_+ \end{bmatrix} = \begin{bmatrix} \vec{x}_0 + \Delta \vec{x} \\ \vec{v}_0 + \Delta \vec{v} \end{bmatrix} = \underbrace{\begin{bmatrix} \mathbb{I} & \mathbf{D}_1 \\ 0 & \mathbb{I} + \Delta t \mathbf{D}_0 \end{bmatrix}}_{\mathbb{M}} \cdot \begin{bmatrix} \vec{x}_0 \\ \vec{v}_0 \end{bmatrix}$$



B-Frame Push in Variable Fields (cont.):

- Push as Explicit Matrix Operator:
(For now Ignoring \vec{E})
- What's Known about \mathbb{M} :
 - Reversible through Time Inversion...
 - \mathbb{M}^{-1} Must Exist
 - \mathbb{M} is a 1:1 Map of $\mathbb{X}_0 \rightarrow \mathbb{X}_+$
 - \mathbb{M} is Unique (Even if \hat{R} was Not...)
 - If \mathbb{M} is Unique, Elements are Unique
 - \mathbb{M} is Constant if \vec{B} is Constant

Hypothesis:

Elements of \mathbb{M} Vary Smoothly, $\vec{x} \in_{\text{Ngbh}} \vec{x}_0$

(Assuming \vec{B} Varies Smoothly with \vec{x})


$$\mathbb{X}_+ = \mathbb{M} \cdot \mathbb{X}_0$$

$$\mathbb{X}_+ = \begin{bmatrix} \vec{x}_+ \\ \vec{v}_+ \end{bmatrix} = \begin{bmatrix} \vec{x}_0 + \Delta \vec{x} \\ \vec{v}_0 + \Delta \vec{v} \end{bmatrix} = \underbrace{\begin{bmatrix} \mathbb{I} & \mathbf{D}_1 \\ 0 & \mathbb{I} + \Delta t \mathbf{D}_0 \end{bmatrix}}_{\mathbb{M}} \cdot \begin{bmatrix} \vec{x}_0 \\ \vec{v}_0 \end{bmatrix}$$



B-Frame Push in Variable Fields (cont.):


- Assuming \vec{B} varies Periodically in ω_c and Linearly in multiples $N \times \omega_c$...





B-Frame Push in Variable Fields (cont.):

- Assuming \vec{B} varies Periodically in ω_c and Linearly in multiples $N \times \omega_c$...






B-Frame Push in Variable Fields (cont.):

- Assuming \vec{B} varies Periodically in ω_c and Linearly in multiples $N \times \omega_c$...?
- Elements of \mathbb{M} take Form:

$$m_{ij} = \bar{m}_{ij} + \frac{n\delta t}{\Delta t} \Delta m_{ij}^{[slow]} + \cos(\omega n \delta t) \Delta m_{ij}^{[c]} + \sin(\omega n \delta t) \Delta m_{ij}^{[s]}$$






B-Frame Push in Variable Fields (cont.):

- Assuming \vec{B} varies Periodically in ω_c and Linearly in multiples $N \times \omega_c$...?
- Elements of \mathbb{M} take Form:

$$m_{ij} = \bar{m}_{ij} + \frac{n\delta t}{\Delta t} \Delta m_{ij}^{[slow]} + \cos(\omega n \delta t) \Delta m_{ij}^{[c]} + \sin(\omega n \delta t) \Delta m_{ij}^{[s]}$$

- Coefficients, Δm_{ij} , can be Sampled using Original Push with $\Delta t = [0, \pi/4, 3\pi/4, 2\pi n]\omega_c$






MULTI-SCALE PARTICLE ADVANCEMENT

B-Frame Push in Variable Fields (cont.):

- Assuming \vec{B} varies Periodically in ω_c and Linearly in multiples $N \times \omega_c$...?
- Elements of \mathbb{M} take Form:

$$m_{ij} = \bar{m}_{ij} + \frac{n\delta t}{\Delta t} \Delta m_{ij}^{[slow]} + \cos(\omega n \delta t) \Delta m_{ij}^{[c]} + \sin(\omega n \delta t) \Delta m_{ij}^{[s]}$$

- Coefficients, Δm_{ij} , can be Sampled using Original Push with $\Delta t = [0, \pi/4, 3\pi/4, 2\pi n]\omega_c$
- Comparison of Fit and Sampled \mathbb{M} from Magnetic Mirror



Closed Circle - Original Push Stepping m_{ij}
Open Circle - Estimated m_{ij} from Sampling



Compounded Push Operator:

- Uniform Field Push can be Subdivided





Compounded Push Operator:

- Uniform Field Push can be Subdivided

$$\begin{aligned}\mathbb{X}(t + \Delta t) &= \mathbb{M}(\Delta t) \cdot \mathbb{X}_0 \\ &= \mathbb{M}^{(p-1)}(\Delta t/p) \cdot \mathbb{X}_0\end{aligned}$$



Compounded Push Operator:

- Uniform Field Push can be Subdivided
- Can be Written as Product of δt Substeps

$$\begin{aligned}\mathbb{X}(t + \Delta t) &= \mathbb{M}(\Delta t) \cdot \mathbb{X}_0 \\ &= \mathbb{M}^{(p-1)}(\Delta t/p) \cdot \mathbb{X}_0 \\ &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]}(\delta t) \cdot \mathbb{X}_0\end{aligned}$$



Compounded Push Operator:

- Uniform Field Push can be Subdivided
- Can be Written as Product of δt Substeps
- Also True if $\mathbb{M}^{[k]}$ varies with t – Equivalent to Explicit Stepping with Smaller Δt

$$\begin{aligned}\mathbb{X}(t + \Delta t) &= \mathbb{M}(\Delta t) \cdot \mathbb{X}_0 \\ &= \mathbb{M}^{(p-1)}(\Delta t/p) \cdot \mathbb{X}_0 \\ &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]}(\delta t) \cdot \mathbb{X}_0\end{aligned}$$



Compounded Push Operator:

- Uniform Field Push can be Subdivided
- Can be Written as Product of δt Substeps
- Also True if $\mathbb{M}^{[k]}$ varies with t – Equivalent to Explicit Stepping with Smaller Δt
- Estimator for $\mathbb{M}^{[k]}$ Reduces Field Evaluations, but not Push Multiplier Operations...

$$\begin{aligned}\mathbb{X}(t + \Delta t) &= \mathbb{M}(\Delta t) \cdot \mathbb{X}_0 \\ &= \mathbb{M}^{(p-1)}(\Delta t/p) \cdot \mathbb{X}_0 \\ &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]}(\delta t) \cdot \mathbb{X}_0\end{aligned}$$



MULTI-SCALE PARTICLE ADVANCEMENT

Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$

$$\tilde{\mathbb{M}} = \prod_{k=0}^{p-1} \mathbb{M}^{[k]}$$



MULTI-SCALE PARTICLE ADVANCEMENT

Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$
- Can be Split into Mean and Deviation

$$\begin{aligned}\tilde{\mathbb{M}} &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]} \\ &= \prod_{k=0}^{p-1} (\bar{\mathbb{M}} + \Delta \mathbb{M}^{[k]})\end{aligned}$$



Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$
- Can be Split into Mean and Deviation
- The Operator is Linearized around $\bar{\mathbb{M}}$

$$\begin{aligned}\tilde{\mathbb{M}} &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]} \\ &= \prod_{k=0}^{p-1} (\bar{\mathbb{M}} + \Delta \mathbb{M}^{[k]}) \\ &= \prod_{k=0}^{p-1} (\mathbb{L} \mathbb{A} \mathbb{R} + \Delta \mathbb{M}^{[k]})\end{aligned}$$



Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$
- Can be Split into Mean and Deviation
- The Operator is Linearized around $\overline{\mathbb{M}}$

$$\begin{aligned}\tilde{\mathbb{M}} &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]} \\ &= \prod_{k=0}^{p-1} (\overline{\mathbb{M}} + \Delta \mathbb{M}^{[k]}) \\ &= \mathbb{L} \left[\prod_{k=0}^{p-1} (\mathbb{A} + \mathbb{R} \Delta \mathbb{M}^{[k]} \mathbb{L}) \right] \mathbb{R}\end{aligned}$$



Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$
- Can be Split into Mean and Deviation
- The Operator is Linearized around $\bar{\mathbb{M}}$
- And Assuming
 $||\Lambda|| \gg ||\mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}||$

$$\begin{aligned}
 \tilde{\mathbb{M}} &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]} \\
 &= \prod_{k=0}^{p-1} (\bar{\mathbb{M}} + \Delta\mathbb{M}^{[k]}) \\
 &= \mathbb{L} \left[\prod_{k=0}^{p-1} (\Lambda + \mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}) \right] \mathbb{R} \\
 &= \mathbb{L} \left[\Lambda^{(p-1)} + \sum_{k=0}^{p-2} (\Lambda^{(p-2)-k} \mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}\Lambda^k) + o(\Delta\mathbb{M}^2) \right] \mathbb{R}
 \end{aligned}$$



MULTI-SCALE PARTICLE ADVANCEMENT

Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbb{M}}...$
- Can be Split into Mean and Deviation
- The Operator is Linearized around $\bar{\mathbb{M}}$
- And Assuming $||\Lambda|| \gg ||\mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}||$
- Summations to Series Solution for Linear and Oscillatory Parts of $\Delta M \rightarrow$ Direct Eval of m_{ij}


$$\begin{aligned}\tilde{\mathbb{M}} &= \prod_{k=0}^{p-1} \mathbb{M}^{[k]} \\ &= \prod_{k=0}^{p-1} (\bar{\mathbb{M}} + \Delta\mathbb{M}^{[k]}) \\ &= \mathbb{L} \left[\prod_{k=0}^{p-1} (\Lambda + \mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}) \right] \mathbb{R} \\ &= \mathbb{L} \left[\Lambda^{(p-1)} + \sum_{k=0}^{p-2} (\Lambda^{(p-2)-k} \mathbb{R}\Delta\mathbb{M}^{[k]}\mathbb{L}\Lambda^k) + o(\Delta\mathbb{M}^2) \right] \mathbb{R}\end{aligned}$$

$$\begin{aligned}m_{ij} &\approx \mathbb{L} \left[\lambda_i^{(p-1)} + \lambda_i^{(p-2)} \mathbb{Q}_{ij} \sum_{k=0}^{p-2} [f_Q(k) \lambda_j^k \lambda_i] \right] \mathbb{R} \\ &\approx \mathbb{L} \left[\lambda_i^{(p-1)} + \lambda_i^{(p-2)} \mathbb{Q}_{ij} \left\{ \begin{array}{l} \sum k \lambda^k, \\ \sum \cos(\alpha k) \lambda^k, \\ \sum \sin(\alpha k) \lambda^k \end{array} \right\} \right] \mathbb{R}\end{aligned}$$



Operation Reduction:

- Given the Gyro-Estimated Compounded Operator $\tilde{\mathbf{M}}...$
- Can be Split into Mean and Deviation
- The Operator is Linearized around $\bar{\mathbf{M}}$
- And Assuming $||\Lambda|| \gg ||\mathbf{R}\Delta\mathbf{M}^{[k]}\mathbf{L}||$
- Summations to Series Solution for Linear and Oscillatory Parts of $\Delta\mathbf{M} \rightarrow$ Direct Eval of m_{ij}



Magnetic Mirror Test



Elementary processes (electron-impact only)


- Collisional excitation/deexcitation



- Collisional ionization/recombination



- Bound-bound radiative transition





Rate Equation

$$K_{ji}^{rad.} = A_{ji}, \quad K_{ji}^{coll.} = N_e \int_{E_{ij}}^{\infty} \sigma_{ij}(\varepsilon) v f(\varepsilon) d\varepsilon$$

$$\frac{dN_i}{dt} = \sum_{R \in R_{xn}} \sum_{j \neq i} (K_{ij}^R N_j - K_{ji}^R N_i)$$


$$dN_i = \sum_{j \neq i} \tilde{\mathbf{K}}_{ij} N_j - \tilde{\mathbf{J}}_i \circ N_i$$

Implicit formulation

$$dN_j = \mathbb{A}_{ij}^{-1} \left[\sum_{j \neq i} \tilde{\mathbf{K}}_{ij} N_j - \tilde{\mathbf{J}}_i \circ N_i \right]$$



LEVEL GROUPING





LEVEL GROUPING


n, m : bin indices; i, j : level indices

Model	Cons. vars.	Size	Lev. Pop.
Full CR	N_i	N_{level}	
Reduced CR (Uniform)	$\Sigma_n = \sum_{i \in n} N_i$	N_{group}	$N_{i \in n} = \frac{g_i \Sigma_n}{\sum_{j \in n} g_j}$
Reduced CR (Boltzmann)	N_{n_0} $\Sigma'_n = \sum_{i \in n'} N_i$	$2 \times N_{\text{group}}$	$N_{i \in n} = \frac{g_i N_{n_0} e^{-\Delta E_i / k T_n}}{g_{n_0}}$ or $N_{i \in n} = \frac{g_i \Sigma'_n e^{-\Delta E_i / k T_n}}{Q'_n}$



ISOTHERMAL HEATING TEST ($T_e = 3 \text{ eV}$)



Comparison of ground state and electron number densities






ISOTHERMAL HEATING TEST ($T_e = 3 \text{ eV}$)

Comparison of excited states number densities






UNIFORM VS. BOLTZMANN





UNIFORM VS. BOLTZMANN






Numerous Previous Merge Methods:



PRIOR MERGING TECHNIQUES

Numerous Previous Merge Methods:

- 2:1 - Cannot Conserve Energy
(Lapenta & Brackbill, JCP 1994)




$$w_m = \sum w_i$$
$$\vec{v}_m = \sum w_i \vec{v}_i / w_m$$
$$w_m v_m^2 < \sum w_i v_i^2 !!!$$



PRIOR MERGING TECHNIQUES

Numerous Previous Merge Methods:

- 2:1 - Cannot Conserve Energy
(Lapenta & Brackbill, JCP 1994)
- Complex Macro-particles with Internal Energy
(Hewett, JCP 2003)




$$w_m = \sum w_i, \quad \vec{v}_m = \sum w_i \vec{v}_i / w_m$$
$$T_m^{(int)} = \left(\sum w_i T_i^{(int)} + \sum w_i v_i^2 - w_m v_m^2 \right) / w_m$$
$$w_m \left(v_m^2 + T_m^{(int)} \right) = \sum \left(w_i v_i^2 + w_i T_i^{(int)} \right) \quad \checkmark$$

Particle Push with $T^{(int)}$?...
Split if $(T^{(int)})^{1/2} \gg v_m$?...
In What Coord-System?...



Numerous Previous Merge Methods:

- 2:1 - Cannot Conserve Energy
(Lapenta & Brackbill, JCP 1994)
- Complex Macro-particles with Internal Energy
(Hewett, JCP 2003)
- Merge to Grid
(Assous et al., JCP 2003, Welch et al., JCP 2007)



Conserve Arbitrary Moments to Grid


Not Explicitly Conserved Lost?
Entropy Generation?
Shape Functions?
Grid Dependence?



Numerous Previous Merge Methods:

- 2:1 - Cannot Conserve Energy
(Lapenta & Brackbill, JCP 1994)
- Complex Macro-particles with Internal Energy
(Hewett, JCP 2003)
- Merge to Grid
(Assous et al., JCP 2003, Welch et al., JCP 2007)

All Introduce Significant Error
and/or Complexity



Conserve Arbitrary Moments to Grid


Not Explicitly Conserved Lost?
Entropy Generation?
Shape Functions?
Grid Dependence?



Merge to Pair → DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

(AFOSR Review 2006)



$$w_m = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \left(\sum_i^{(n+2)} w_i \vec{v}_i \right) / w_m$$

$$\overline{v^2} = \left(\sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2 \right) / w_m$$

$$w_{(a/b)} = w_m / 2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{\mathcal{R}} \sqrt{\overline{v^2}}$$

(Similarly: $\vec{x}_{(a/b)} = \vec{x} \pm \hat{\mathcal{R}} \sqrt{\overline{x^2}}$)



CONSERVATIVE MERGE


Merge to Pair \rightarrow DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

(AFOSR Review 2006)

Selection of Near Neighbors in VDF Limits Thermalization

(\approx Near Neighbor Pairs in 2:1 Merges that Limit Numerical Cooling)



$$w_m = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \left(\sum_i^{(n+2)} w_i \vec{v}_i \right) / w_m$$

$$\overline{v^2} = \left(\sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2 \right) / w_m$$

$$w_{(a/b)} = w_m / 2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{R} \sqrt{\overline{v^2}}$$

(Similarly: $\vec{x}_{(a/b)} = \vec{x} \pm \hat{R} \sqrt{\overline{x^2}}$)



Phase-Space Decomposition

- Given a Set of Particles...






OCTREE BINNING

Phase-Space Decomposition


- Given a Set of Particles...
- Particles Binned in Octants





Phase-Space Decomposition

- Given a Set of Particles...
- Particles Binned in Octants
- Octants Recursively Sub-Divided






OCTREE BINNING

Phase-Space Decomposition

- Given a Set of Particles...
- Particles Binned in Octants
- Octants Recursively Sub-Divided
- Recursion Halted at 1-Particle/Bin or Other Criteria such as Bin-Density






OCTREE BINNING

Phase-Space Decomposition

- Given a Set of Particles...
- Particles Binned in Octants
- Octants Recursively Sub-Divided
- Recursion Halted at 1-Particle/Bin or Other Criteria such as Bin-Density


Restricts Phase-Space Diffusion to
Within Local Bins





0D-MERGE EXAMPLES


Comparison of Random vs. Octree Merge Partner Selection (Note: Mass, Momentum, and Kinetic Energy Both Exactly Conserved)





0D-MERGE EXAMPLES

Comparison of Random vs. Octree Merge Partner Selection (Note: Mass, Momentum, and Kinetic Energy Both Exactly Conserved)






BEAM IN POTENTIAL WELL

Collisionless Particles in Well

- 6000 Unmerged Particles
- Reproduces 3-4 Orders of Magnitude






BEAM IN POTENTIAL WELL

Collisionless Particles in Well

- 6000 Unmerged Particles
- Reproduces 3-4 Orders of Magnitude
- Random Merge -> Thermalization
- 3000 First Point, 1500 First Cross
- Bi-Maxwellian Specifically Difficult






BEAM IN POTENTIAL WELL

Collisionless Particles in Well

- 6000 Unmerged Particles
- Reproduces 3-4 Orders of Magnitude
- Random Merge -> Thermalization
- 3000 First Point, 1500 First Cross
- Bi-Maxwellian Specifically Difficult
- Octree Merge Significantly Better






BEAM IN POTENTIAL WELL

Collisionless Particles in Well

- 6000 Unmerged Particles
- Reproduces 3-4 Orders of Magnitude
- Random Merge -> Thermalization
- 3000 First Point, 1500 First Cross
- Bi-Maxwellian Specifically Difficult
- Octree Merge Significantly Better
- Merge & Split Adapts Particle Count






BEAM IN POTENTIAL WELL

Collisionless Particles in Well


- 6000 Unmerged Particles
- Reproduces 3-4 Orders of Magnitude
- Random Merge -> Thermalization
- 3000 First Point, 1500 First Cross
- Bi-Maxwellian Specifically Difficult
- Octree Merge Significantly Better
- Merge & Split Adapts Particle Count
- Computational Particles per Cell Vastly Different






DC-Diode Test Case:

- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot



Control




Merged




DC-BREAKDOWN

DC-Diode Test Case:

- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode \rightarrow Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode



Control




Merged



DC-Diode Test Case:


- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode \rightarrow Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode
- Chain-Branching Needs Merge





DC-Diode Test Case:


- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode \rightarrow Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode
- Chain-Branching Needs Merge
- Merge Overhead Rapidly Negligible





DC-Diode Test Case:

- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode → Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode
- Chain-Branching Needs Merge
- Merge Overhead Rapidly Negligible
- Control: Parts/Cell \propto Density





Control

Merged



DC-Diode Test Case:

- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode \rightarrow Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode
- Chain-Branching Needs Merge
- Merge Overhead Rapidly Negligible
- Control: Parts/Cell \propto Density
- Merge: Parts/Cell Much Reduced







DC-BREAKDOWN

DC-Diode Test Case:

- Full 3D Electrostatic-PIC
- Averaged to 1D XT-Plot
- 250V Cathode \rightarrow Anode
- MCC-Ionization Collisions
- Secondary Emission at Cathode
- Chain-Branching Needs Merge
- Merge Overhead Rapidly Negligible
- Control: Parts/Cell \propto Density
- Merge: Parts/Cell Much Reduced
- Despite Identical Densities



Control




Merged



Integrate R&D w/ Production TODO:

- High-Order Fluid/MHD GPU Models
(Le/Cole*/Bilyeu PhD Research)
- GPU Accelerated Chemical Kinetics /
CR Ar-Ne-Xe-Molecular Models
(Le/Cole*/Kapper* PhD Research)
- Phase-Space Reconstruction/Vlasov
(Martin/Bilyeu/TBD)
- Implicit / Multiscale GPU-Accelerated
PIC
(Lederman/Gimelsheins/Martin/TBD)

*Note: Former Co-op Student Work to be Integrated into Framework





END



Thank You

Questions?